

Facility QI / QM History - Chronic Care

Domain - QI / QM	Jun'02 Facility Percent	Sep'02 Facility Percent	Dec'02 Facility Percent	Mar'03 Facility Percent	Jun'03 Facility Percent	Jun'03 Comparison Percent
Accidents						
1.1 Incidence of new fractures	0.0	1.6	2.1	1.0	1.1	1.7
1.2 Prevalence of falls	17.8	19.6	20.2	19.5	18.3	13.9
Behavior/Emotional Patterns						
2.1 Residents who have become more depressed or anxious	0.0	24.6	17.4	25.0	18.4	15.4
2.2 Prevalence of behavior symptoms affecting others	25.7	21.5	17.8	15.9	15.4	18.7
High risk	30.0	24.3	20.2	20.3	20.3	22.0
Low risk	19.5	16.7	12.5	7.7	5.7	8.4
2.3 Prevalence of symptoms of depression without antidepressant therapy	7.9	5.5	3.1	3.5	3.8	4.5
Clinical Management						
3.1 Use of nine or more different medications	38.6	39.9	44.2	43.4	40.4	54.9
Cognitive Patterns						
4.1 Incidence of cognitive impairment	0.0	10.0	3.0	9.8	13.2	11.9
Elimination/Incontinence						
5.1 Low-risk residents who lost control of their bowels or bladder	37.0	36.9	32.5	31.9	36.8	45.5
5.2 Residents who have/had a catheter inserted and left in their bladder	2.0	4.3	8.5	8.8	5.8	6.2
High risk	0.0	0.0	11.1	15.8	11.1	11.7
Low risk	2.0	4.5	8.1	7.4	4.7	5.2
5.3 Prevalence of bladder or bowel incontinence without a toileting plan	14.6	17.6	16.4	20.0	21.3	31.1
5.4 Prevalence of fecal impaction	1.0	0.6	0.0	0.9	1.0	0.1
Infection Control						
6.1 Residents with a urinary tract infection	12.9	9.8	8.5	15.0	15.4	6.8
Nutrition/Eating						
7.1 Residents who lose too much weight	9.0	8.8	8.8	10.1	9.9	9.0
7.2 Prevalence of tube feeding	0.0	1.2	3.1	3.5	1.9	4.1
7.3 Prevalence of dehydration	0.0	0.0	0.0	0.0	0.0	0.3
Pain Management						
8.1 Residents who have moderate to severe pain	15.8	8.0	7.0	15.0	11.5	9.4
High risk	0.0	17.2	10.4	22.2	21.6	14.2
Low risk	15.8	6.0	4.9	8.5	1.9	6.3
Physical Functioning						
9.1 Residents whose need for help with daily activities has increased	0.0	16.4	15.1	13.7	12.2	15.4
9.2 Residents who spend most of their time in a bed or chair	3.0	6.7	10.1	8.0	4.8	2.4
9.3 Residents whose ability to move in and around their room got worse	0.0	19.1	13.0	12.7	12.2	15.0
High risk	0.0	12.9	14.8	14.3	14.8	17.2
Low risk	0.0	31.3	8.7	8.7	5.0	9.4
9.4 Incidence of decline in ROM	0.0	29.2	43.1	61.6	61.2	8.7
Psychotropic Drug Use						
10.1 Prevalence of antipsychotic use in the absence of psychotic conditions	17.0	16.7	16.5	16.1	14.4	18.1
High risk	22.2	52.0	61.1	40.0	28.6	39.8
Low risk	15.9	10.2	9.2	12.4	12.2	14.2
10.2 Prevalence of antianxiety/hypnotic use	23.0	18.5	17.3	17.9	18.3	14.3
10.3 Prevalence of hypnotic use more than two times in last week	7.9	7.4	5.4	6.2	5.8	2.4
Quality of Life						
11.1 Residents who were physically restrained	14.9	11.0	5.4	4.4	4.8	4.6
11.2 Prevalence of little or no activity	26.7	28.8	36.4	45.1	41.3	7.6
Skin Care						
12.0 Residents with pressure ulcers	5.9	8.6	10.9	13.3	9.6	8.8
High risk	9.4	14.1	16.9	18.2	12.5	13.1
Low risk	0.0	1.4	3.4	8.6	7.1	3.4